



ZIERING MEDICAL

EYEBROW AWAKENINGS

DAY OF SURGERY:

- After dinner, start taking medications one hour apart from each other with food.
- Start icing underneath eyebrows (on eyelids)
- Remove all dressings upon arrival home the night of your surgery
- Use button down shirt
- Sleep at a 45-degree angle

REUSABLE ICE PACK

To be used to reduce post-operative swelling. Do NOT apply directly to eyebrows.

EYE THERAPY

To be used to minimize bruising and swelling around the eye area

POST-OPERATIVE COMFORT PILLOW

To be used for 3-4 nights following surgery. It helps to stabilize the head and neck while resting or sleeping to minimize movement for protection of grafts.

HERBAL SKIN CLEANSER

This is an alcohol-free cleanser that is used to wash the face. It is also used to create the cleanser mix to wash the eyebrows. Follow the instructions given on the post-operative care sheet when washing the eyebrows.

CLEANSER MIX

Herbal skin cleanser diluted with saline or distilled water to spritz eyebrows for cleansing.

SALINE SOLUTION/DISTILLED WATER

Spritz saline or distilled water to rinse off the cleanser mix, then pat dry. Do NOT rub.

BACITRACIN OINTMENT

To be used on the donor area to prevent an infection

FOLICAN GRAFT ENHANCEMENT SPRAY

This spray manages crusting and optimizes the healing process by using the active ingredient copper tripeptide-1.



ZIERING MEDICAL

SIX POST-OPERATIVE LASER LIGHT THERAPY TREATMENTS

These treatments will shorten the healing process as well as reduce swelling and overall discomfort after your procedure.

BIO OIL

Apply to suture line daily starting 1 week after suture removal. (See Bio Oil instruction sheet)

Disregard if you had FUE procedure

GROOMING KIT

Scissors

These scissors are used for constant maintenance of trimming your eyebrows. You may begin to use 10 days after procedure.

Eyebrow brush with comb

This brush is used for daily grooming of your eyebrows. You may begin to use 10 days after your procedure.

Eyebrow gel

Brow gel is used to hold the eyebrows in place as it grooms, sets and defines. You may begin to use 1 month after your procedure.